

FERTILIZERS TREATMENTS IN CASE OF EMERGENCY

BACTIBAC

SIMPLE MINERAL FERTILIZER SOLUTION OF POTASSIUM SALTS BTC

COMPOSITION

POTASSIUM OXIDE (K₂O) water soluble: 10%

FEATURES

BACTIBAC can be used on a wide range of plant species (all crops, edible and ornamental plants). Potash is present in a particular form that allows BACTIBAC to be particularly useful for the prevention of many physiopathologies. It improves the natural strength of plant organs, the epidermal pigmentation, the endogenous response to pest attacks and it finally facilitates the cold storage.

DOSES AND USE PATTERNS

We recommend at least two annual treatments: the first 20-30 days after the germination for deciduous woody plants and shrubs and the latter close to the natural fall of leaves in autumn. Concerning evergreen woody plants and shrubs we recommend two treatments: the first in spring when temperatures increase and the latter in autumn.

Herbaceous plants can be treated only one time per production cycle if it does not exceed 30-40 days, or more frequently for longer production cycle.

DOSES

Foliar treatments:

Herbaceous plants: 1,5-2,5 kg per hectare

Woody plants and shrubs: 3-4 kg per hectare

Floriculture: 0,5-0,8 kg per hectare (depending of the species)

PACKING: 1 KG BOTTLES-BOXES OF 20 PIECES, 5 KG CANS.

